

HEPATITIS B VACCINE

Hepatitis B is a serious disease.

The hepatitis B virus (HBV) can cause short-term (acute) illness that leads to:

- loss of appetite
- diarrhea and vomiting
- tiredness
- jaundice (yellow skin or eyes)
- pain in muscles, joints, and stomach

It can also cause long-term (chronic) illness that leads to:

- liver damage (cirrhosis)
- liver cancer
- death

About 1.25 million people in the U.S. have chronic HBV infection.

Each year it is estimated that:

- 80,000 people, mostly young adults, get infected with HBV
- More than 11,000 people have to stay in the hospital because of hepatitis B
- 4,000 to 5,000 people die from chronic hepatitis B

Hepatitis B vaccine can prevent hepatitis B. It is the first anti-cancer vaccine because it can prevent a form of liver cancer.

Hepatitis B virus is spread through contact with the blood and body fluids of an infected person. A person can get infected in several ways, such as:

- by having unprotected sex with an infected person
- by sharing needles when injecting illegal drugs
- by being stuck with a used needle on the job
- during birth when the virus passes from an infected mother to her baby

About 1/3 of people who are infected with hepatitis B in the United States don't know how they got it.

- 1) Everyone 18 years of age and younger
- 2) Adults over 18 who are at risk

Adults at risk for HBV infection include:

- people who have more than one sex partner in 6 months
- men who have sex with other men
- sex contacts of infected people
- people who inject illegal drugs
- health care and public safety workers who might be exposed to infected blood or body fluids
- household contacts of persons with chronic HBV infection
- hemodialysis patients

If you are not sure whether you are at risk, ask your doctor or nurse.

D People should get 3 doses of hepatitis B vaccine according to the following schedule. *If you miss a dose or get behind schedule, get the next dose as soon as you can. There is no need to start over.*

- The second dose must be given at least 1 month after the first dose.
overt⁷



In the rare event that you or your child has a serious reaction to a vaccine, a federal program has been created to help you pay for the care of those who have been harmed.

For details about the National Vaccine Injury Compensation Program, call **1-800-338-2382** or visit the program's website at **www.hrsa.gov/osp/vicp**

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department's

What should I look for?

Any unusual condition, such as a serious allergic reaction, high fever or unusual behavior. Serious allergic